

FODDER CROPS

FRAMEWORK PROGRAM FOR MINERAL™ USE ON FODDER CROPS Soya, Fodder peas					
Development phase	Measure	Mineral™	Solution		
before sowing	Soaking seeds in a solution for 20 minutes	MINERAL [™] YELLOW	2 %		
	- speeds up germination		(20 ml / 1 l water) (2 dcl / 10 l water) (2 l / 100 l water)		
immediately after sowing	Watering	MINERAL™ GREEN	4 %		
	if the technology of production permits soak the soil with 10 000 l water/ha before watering with Mineral™ Green		20 / 500 water / ha		
	- delivers nutrients and water into the soil - speeds up germination				
	 It is not appropriate on sandy soils in intensive production use disinfected seeds 				

14 days after the emergence	Spraying - prevention	MINERAL™ YELLOW	4 %
	Repeat regularly on all above ground parts of plants, blossoms as well, once in 14 days		10 / 250 water / ha
	 increases the resistance of plants against diseases and stress foliar feeding (through leaves) increases quality of the crop 		
15 days after the emergence	Spraying - prevention	MINERAL™ RED	4 %
	Repeat regularly on all above ground parts of plants, blossoms as well, once in14 days		10 / 250 water / ha
	Always a day after spraying with Mineral™ Yellow		
	 increases the resistance of plants against pests and stress foliar feeding (through leaves) 		
IN THE EVENT OF			
the appearance of first signs of a disease	Spraying plants on all above ground parts of plants - curative	MINERAL™ YELLOW	7 %
(mold etc)	Repeat at least 3 times every other day (day 1., 3., 5.)		17,5 l / 250 l water / ha
	- market growers should use an appropriate fungicide		

the appearance of first signs of a pest	Spraying plants on all above ground parts of plants - curative	MINERAL™ RED	7% 17,51 /
(aphids, mites, etc)	Repeat at least 3 times every other day (day 1., 3., 5.)		250 l water / ha
	Do not spray at the same time with Mineral [™] Yellow.		
	- market growers should use an appropriate insecticide		